

Policy & Procedure (P&P)

Policy Title :

Anti-A1 Typing

Department	Index No.	Scope
Laboratory & Blood Bank	LAB-063	All Laboratory staff
Issue Date	Revision NO	Effective Date
1437/06/12	3	1440/07/23
Review Due Date	Related Standard NO.	Page Number#
1442/07/23	CBAHI (LB. 50)	3

01. Policy :

01.1. The blood bank confirms all specimens suspected to be A2 or A2B blood sub-groups using the anti-A1 lectin.

02. Definition :

02.1. N/A

03. Purpose :

03.1. All specimens where A2 or A2B blood group is suspected is confirmed with anti-A1 lectin.

04. Procedure :

04.1. Specimen Requirements

Blood collected with anticoagulant is used.

Specimens collected in CPD, CPDA-1, sodium citrate, oxalate, EDTA, or heparin are all acceptable.

Testing should be performed as soon as possible.

If a delay in testing occurs, specimens should be stored at 2-8 degrees centigrade and tested within 48 hours.

No patient preparation is required.

04.2. Materials

04.2.1. Glass test tubes

04.2.2. Blood Bank pipettes

04.2.3. Isotonic saline 0.85% NaCl

04.2.4. Calibrated serologic centrifuge

04.2.5. Microscope and slides

04.3. Reagents

04.3.1. Anti-A1 Lectin

04.4. Method

04.4.1. Wash an appropriate volume of patient red blood cells two times with isotonic saline.

04.4.2. Prepare a 3% to 5% suspension of red blood cells in isotonic saline.

04.4.3. To a labeled test tube add one drop of anti-A1 lectin.

04.4.4. Using a plastic pipette add one drop of the patient's cell suspension

04.4.5. Mix well and centrifuge according to the optimum saline spin time listed for the centrifuge.

04.4.6. Resuspend the cells by gentle agitation and examine macroscopically and microscopically.

04.4.7. Record results on worksheet.

04.5. Quality Control

Reagents are to be tested with appropriate positive and negative controls daily and the results recorded. See "Daily Quality Control" for procedure.

04.6. Reporting Results:

04.6.1. Interpretation

Agglutination of the red cells is a positive test result and indicates the presence of A1 antigen.

No agglutination is a negative test result and indicates the absence of the A1 antigen and the blood group is A2 or A2B.

Reactions should show the following pattern with group A red blood cells.

Anti-A1 Lectin	Interpretation
+	A1
-	A2

04.7. Limitations:

04.7.1. Testing must always be performed at room temperature (20 to 30 C) and never incubated at higher

temperatures.

04.7.2. The A1 antigen is not fully expressed on the red blood cells of newborn infants and false negative results may occur

04.7.3. Leukemia may produce a weekend form of A.

05. Responsibilities :

05.1. All laboratory of Al-Qunfudah General Hospital.

06. Equipment & Forms

06.1. Blood Grouping Book






07. Attachment :

07.1. N/A

08. Reference

08.1. The Technical manual of the American Association of Blood Banks.

Preparation , Reviewing & Approval Box

	NAME	POSITION	SIGN & STAMP	DATE
Prepared By	Dr RAJA NACER SASSI	Head of Blood Bank		
Reviewed By	Mr. ABDULHADI ASHIRI	Lab & B.Bank HOD		
Document Reviewed By	Ms. SADIAH ALMAHMOUDI	TQM Director		٢٠/٥/٢٠٢٠
Reviewed By	Dr. AGEEL ALGANIMI	Medical Director		
Approved By	Dr. ABDULLAH ALJABRI	Hospital Director		١٤٤٠/٧/١٧

